

279-3289

9/26/2014

Page 1 of 2

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460



OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

September 26, 2014

Mr. Evan M. Parenti
Senior Registration Specialist
FMC Corporation
1735 Market Street
Philadelphia, PA 19103

Subject: Label Amendment – Adding General Control Claims for Ticks & Use Direction
for Residential Gardens.
Product Name: Zeta-Cyp 1.5 EW OTC Insecticide
EPA Registration Number: 279-3289
Application Date: April 14, 2014
Decision Number: 490011

Dear Mr. Parenti:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). If you have any questions, please contact BeWanda Alexander by phone at (703)305-7460, or via email at alexander.bewanda@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "Richard Gebken".

Richard Gebken, Product Manager 10
Insecticide Branch
Registration Division (7505P)
Office of Pesticide Programs

278

ZETA-CYP 1.5 EW

OTC Insecticide — For Homeowner Use

For Outdoor Residential Homeowner Use Only.

EPA Reg. No. 279-3289	EPA Est. 279-	
Active Ingredient:		By Wt.
Zeta-Cypermethrin*	17.1%	
Other Ingredients:	82.9%	
	100.0%	

Zeta-Cyp 1.5 EW OTC Insecticide contains 1.5 pounds of active ingredient per gallon.


*Cis/trans ratio: Max. 55%(±) cis and min. 45% (±) trans.

KEEP OUT OF REACH OF CHILDREN

WARNING

STOP – Read the label before use.

See other panels for additional precautionary information.



FIRST AID	
If Swallowed	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
If on Skin or Clothing	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If in Eyes	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If Inhaled	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-331-3148 for emergency medical treatment information.	
NOTE TO PHYSICIAN	
Skin contact (vapor or spray) may be followed by transient tingling or numbness, usually of the face, but this subsides without treatment. This product is a pyrethroid. If large amounts have been ingested, the stomach and intestines should be evacuated. Treatment is symptomatic and supportive.	
For Information Regarding the Use of this Product Call 1-800-321-1FMC (1362).	

PRECAUTIONARY STATEMENTS

Hazards to Humans (and Domestic Animals)

Warning

May be fatal if swallowed. Harmful if inhaled or absorbed through the skin. Causes moderate eye irritation. Avoid breathing vapors or spray mist. Avoid contact with skin, eyes or clothing. Wear long-sleeved shirt, long pants, shoes, socks, gloves and protective eyewear. Remove and wash contaminated clothing before reuse. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Get medical attention if irritation persists.



FMC Corporation
Agricultural Products Group
1735 Market Street
Philadelphia, PA 19103

NET CONTENTS:

Environmental Hazards

This product is extremely toxic to fish and aquatic invertebrates. Do not apply directly to water. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems. Do not contaminate water when disposing of equipment washwaters or rinsate.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops if bees are visiting the treatment area.

{Note to Reviewer:

All optional text enclosed in [brackets].

All notes to reviewer appear in bracketed {italics}.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Before Use, Read and follow the directions below:

SHAKE WELL BEFORE USING

STORAGE AND DISPOSAL

Prohibitions: Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Keep from freezing. Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. For rigid containers: After partial use, replace lids and close tightly.

Product Disposal & Container Handling:

Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

(For containers less than 5 gallons): Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available.

(For containers more than 5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse: Empty the remaining contents into a storage container or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over storage container or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

(1) An appropriate solvent would include a solvent that will dissolve technical material. The rinsate may be used later to formulate a product or may be disposed in accordance with local, state and national environmental laws, rules, standards and regulations.

Returnable/Refillable Containers: Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

{Product Descriptions}

{Option 1}

{Farmstead Insecticide}

[Zeta-Cyp 1.5 EW OTC Insecticide is formulated to kill the listed [insects][pests] that pose harm and nuisance to country and suburban homes and farmsteads. Zeta-Cyp 1.5 EW OTC Insecticide is a versatile tool against insects that can infest and attack lawns, vegetable and flower gardens. Ideal for use with all standard spray equipment including hose-end, pump-style, 12-volt spot, ATV and tow-behind sprayers.]

{Option 2}

{Lawn & Perimeter Insect Control}

[Zeta-Cyp 1.5 EW OTC Insecticide is an insecticide that's formulated to protect country and suburban lawns, homes and structures by quickly killing listed outdoor [insects][pests] by contact—before they can have a chance to enter homes and become a nuisance. Used as a [perimeter], [broadcast], [vegetable garden], [flower garden] spray, Zeta-Cyp 1.5 EW OTC Insecticide can provide up to [4] [6] [8] [12] [weeks] [2] [3] [months]* of continuous protection against listed [insects][pests][over XX lawn and home invading [insects][pests]]. Ideal for use with all standard spray equipment including hose-end, pump-style, 12-volt spot, ATV and tow-behind sprayers.]

*except fleas

{Option 3}

{Lawn & Garden Broadcast Spray}

[Zeta-Cyp 1.5 EW OTC Insecticide is formulated to give you protection against listed outdoor lawn, garden and home-invading [insects][pests]. Whether it's stopping the kinds of outdoor insects that damage lawns and plants or stopping home-invading insects outside, Zeta-Cyp 1.5 EW OTC Insecticide was made to give you protection for up to [4] [6] [8] [12] [weeks] [2] [3] [months]*. Ideal for use with all hose end and pump-style sprayers.]

*except fleas

{Option 4}

[Zeta-Cyp 1.5 EW OTC Insecticide is an insecticide that's formulated to protect homes by killing listed insects outdoors—before they can have a chance to enter homes and become a nuisance. Used as a perimeter [broadcast, vegetable garden, flower garden] spray around homes, buildings and structures, Zeta-Cyp 1.5 EW OTC Insecticide can provide up to [4] [6] [8] [12] [weeks] [2] [3] [months]* [of] continuous protection against home invading insects.]

*except fleas

{OPTIONAL MARKETING CLAIMS}

{General Use Claims}

- Easy-to-use.
- Kills Ants, Earwigs, Silverfish & Other Listed Pests
- Kills by contact.
- Kills [over] 90+ listed insects [by contact].
- Quickly kills listed insects.
- Quickly kills listed insects [by contact].
- Prevents new infestations.
- Fast-acting formula.
- Long lasting* formula.
- *For up to 3 months. Except fleas.
- Protects [for] up to [4] [6] [8] [12] [weeks] [2] [3] [months] *.
- *except fleas
- Keeps protecting up to [2] [3] months*.
- *except fleas
- Keeps killing [for] up to [2] [3] months*.
- *except fleas
- Up to [2] [3] months protection*.
- *except fleas

- Up to [4] [6] [8] [12] [weeks] [2] [3] [months] continuous protection*.
- *except fleas
- Protects lawns, vegetables, flowers, trees and shrubs.
- Kills over 90 listed [insects][pests] in lawns and gardens.
- For all 12-volt sprayers. Instructions inside.
- Ideal for all 12-volt spray equipment.
- For Country and Suburban Lawns, Homes & Structures
- Treats XXX sq. ft. of lawn
- Makes XXX gals of Spray
- 25% More Free! Bonus Size VS 32 oz. size
- 25% More* *VS 32 oz. size
- For hose-end and tank sprayers. Instructions inside.
- Kills Over [xxx] [500] listed insects [by contact].
- Kills Over [xxx] [500] listed insects in lawns and gardens.
- Creates a continuous killing zone around [foundation[s]] [walls]

*except fleas

MIXING INSTRUCTIONS

When using a hose-end sprayer*

Zeta-Cyp 1.5 EW OTC Insecticide may be applied with a hose-end sprayer designed to be attached to a garden hose. Such a sprayer is pre-set to deliver 2.4 fluid ounces of concentrated product per gallon of water [equivalent to a 0.56% solution]. Alternatively, a hose-end sprayer attachment, which can be adjusted to the proper settings, may be used.

With the water turned off at the faucet, attach a garden hose to the sprayer adapter. Make sure the hose is firmly attached and the sprayer is in the off position. Turn the water supply on at the faucet. Water pressure will build in the hose. To apply the product, turn the sprayer control to the on position. For best results, hold the sprayer and plastic container level and direct the sprayer in a sweeping side-to-side movement on lawns, landscaping and shrubs. Water and this product are now being mixed at the proper rate. One gallon of diluted product (For Perimeter: 2.4 fluid ounces of concentrate) (For Lawn & Ornamental Garden: 1.6 fluid ounces of concentrate) will cover 4,000 square feet.

When using 12 Volt/Tow-Behind Boom Sprayers†

To calibrate your application equipment, consult your equipment owner's manual for speed and pressure settings required to deliver 0.5 to 1 gallon of spray solutions to treat 1,000 sq ft. For example, at 4 mph a sprayer might deliver ½ gallon per 1,000 sq ft at 40 psi.

To apply evenly, drive at a constant speed that delivers the prescribed amount of spray required for the area to be treated.

Mix the appropriate amount of product and water in the sprayer using the following table as a guide:

Table 1: 12 Volt and Pull Behind Sprayers Application Guide

Tank Size	Spray Volume	Coverage Area (sq ft)	Concentrate to add to tank (0.1fl. oz rate)	Concentrate to add to tank (0.4 fl. oz rate)	Concentrate to add to tank (0.6 fl. oz rate)
15 gal	0.5 gal/100 sq ft	30,000	3 fl oz	12 fl oz	18 fl oz
	1 gal/100 sq ft	15,000	1.5 fl oz	6 fl oz	9 fl oz
	5 gal/100 sq ft	3,000	0.3 fl oz	1.2 fl oz	1.8 fl oz
25 gal	0.5 gal/100 sq ft	50,000	5 fl oz	20 fl oz	30 fl oz
	1 gal/100 sq ft	25,000	2.5 fl oz	10 fl oz	15 fl oz
	5 gal/100 sq ft	5,000	0.5 fl oz	2.0 fl oz	3.0 fl oz

*Spray method for use only with perimeter treatment or lawn and ornamental gardens.

†Spray method for use only in pastures, perimeter treatment or home lawn and ornamental gardens.

1. HOME INVADING PESTS OUTDOORS

Pests: Ants, boxelder bugs, chiggers, crickets, earwigs, elm leaf beetles, fire-brats, fleas (adults), flies (excluding biting flies), millipedes, blacklegged and brown dog ticks, silverfish & sowbugs (pillbugs).

PERIMETER TREATMENT:

How much to Apply:

When using a tank sprayer

Use 0.60 fl oz of this product per gallon of water to be sprayed over 1,000 sq ft.

Mixing Instructions			
	For Spot Applications	For Broadcast Applications	
	1 Gallon	15 Gallons	25 Gallons
Amount of Water in Sprayer			
Amount of Product to add to sprayer	0.6 fl oz (3.6 tsp)	9 fl oz (18 Tbsp)	15 fl oz (30 Tbsp)
Area to Treat in Square Feet	1,000	15,000	25,000

Do not use household utensils to measure Zeta-Cyp 1.5 EW OTC Insecticide. Utensils such as spoons and measuring cups must not be used for food purposes after use in measuring pesticides.

How to Apply:

Close doors and windows prior to treatment. Use a hose-end or tank sprayer to spray a 5 to 10 foot band of soil around the [house][structure] next to the foundation. Spray the foundation of the [house][structure] to a height of 2 to 3 feet. House siding may also be spot treated if pests such as ants, boxelder bugs, chiggers, crickets, earwigs, elm leaf beetles, firebrats, flies (excluding biting flies), millipedes, silverfish & sowbugs (pillbugs) are present.

Aim spray at nest openings in ground, bushes, and in cracks and crevices which may harbor nests, saturating nest openings and contacting as many insects as possible.

People and pets may re-enter the area once spray has dried.

Where to Use:

Use Zeta-Cyp 1.5 EW OTC Insecticide outdoors as a perimeter spray to obtain up to 3 months* of continuous protection around [homes.] [buildings.] [stables.] [barns] [and] [other structures]. Spray 5 to 10 feet outwards from structures and 2 to 3 feet up the foundation to kill insect pests and provide up to 3 months* of continuous protection.

*except fleas

When to Use:

Application should be made in the late evening when insects are at rest.

[Coverage: One bottle, treats XXX A or XX sq ft]

2. LAWNS AND ORNAMENTAL GARDENS

Pests: Ants, armyworms, annual bluegrass weevil, billbugs, black turfgrass Ataenius, centipedes, chinch bugs, crickets, cutworms, earwigs, fleas (adults), grasshoppers, Japanese beetles, leafhoppers, mealybugs, millipedes, mole crickets, pillbugs, sod webworms, sow bugs and blacklegged and brown dog ticks.

Use higher volume applications if the thatch layer is excessive or if a relatively long mowing height is being maintained.

Where to Use: Use Zeta-Cyp 1.5 EW OTC Insecticide on lawns, vacant lots, ornamental trees, shrubs, and flowers and other similar areas where insect pests are present.

When to Use: Treat lawns and vegetation when insects or damage first appears. Retreatment may be necessary during periods of high pest pressure. Repeat application is necessary only if there are signs of renewed insect activity.

How Much to Apply:

Mix 0.1 – 0.4 fl oz in 1 gallon of water to treat 1,000 sq ft.

For control of fleas (adults) and blacklegged and brown dog ticks use 0.4 fl oz in 1 gallon of water to treat 1,000 sq ft.

Mixing Instructions				
Tank Size	Spray Volume	Coverage Area (sq ft)	Concentrate to add to tank (0.1fl. oz rate)	Concentrate to add to tank (0.4 fl. oz rate)
1 gal	1 gal/1000 sq ft	1,000	0.1 fl oz (0.6 tsp)	0.4 fl oz (2.4 tsp)
	5 gal/1000 sq ft	3,000	0.3 fl oz (1.8 tsp)	1.2 fl oz (2.4 Tbsp)
15 gal	1 gal/1000 sq ft	15,000	1.5 fl oz (3 Tbsp)	6 fl oz (12 Tbsp)
	5 gal/1000 sq ft	3,000	0.3 fl oz (1.8 tsp)	1.2 fl oz (2.4 Tbsp)
25 gal	1 gal/1000 sq ft	25,000	2.5 fl oz (5 Tbsp)	10 fl oz (20 Tbsp)
	5 gal/1000 sq ft	5,000	0.5 fl oz (1 Tbsp)	2.0 fl oz (4 Tbsp)

Do not use household utensils to measure Zeta-Cyp 1.5 EW OTC Insecticide. Utensils such as spoons and measuring cups must not be used for food purposes after use in measuring pesticides.

How to Apply:

Apply Zeta-Cyp 1.5 EW OTC Insecticide as a broadcast treatment. Use application volumes of up to 5 gallons per 1000 sq ft to get uniform coverage when treating dense grass foliage. Lawn should not be longer than 3 inches at the time of application. Arid climates generally require the higher volume. Use higher volume applications if the thatch layer is excessive or if a relatively long mowing height is being maintained.

People and pets may re-enter the area once spray has dried.

[Coverage: One bottle, treats XXX A or XX sq ft]

3. HOME VEGETABLE GARDENS

[Kills: [Aphids.] [Armyworms.] [Corn earworm.] [Cucumber beetles.] [Cutworm.] [Imported Cabbageworm Leafhoppers.] [Crickets.] [Grasshoppers.] [Ground Beetles.] [Leafminers (adults).] [Lygus Bugs.] [Stinkbugs.] [Whiteflies.] [Wireworm (adults)] and XX other listed pests.]

Where to Use:

Home vegetable gardens with brassica (cole), leafy vegetables, bulb vegetables, fruiting vegetables, root and tuber vegetables and sweet corn.

When to Use:

Treat vegetable gardens when insects or damage first appear. Retreatment may be necessary during periods of high pest pressure. Repeat application is necessary only if there are signs of renewed insect activity.

How to Apply:

Apply as insects appear using sufficient water to obtain full coverage of foliage.

Do not exceed maximum allowable rate. Use of a hose-end sprayer is not recommended.

[Coverage: One bottle treats XXX A or XX sq ft]

How Much to Apply:

[See the below tables for these specific rates and amounts of water to apply.]

Application Rate	Per 1,000 sq ft	Per 5,000 sq ft	Per 10,000 sq ft
		0.1 fl oz (2.96ml) (0.6 tsp)	0.5 fl oz (14.8 ml) (3.0 tsp)
Method of Application	<p>Apply as insects appear using sufficient water to obtain full coverage of foliage (minimum of ¼ gallon per 1000 sq ft).</p> <p>Repeat applications as necessary to maintain control.</p> <p>Use lower rates under light to moderate insect pressure. Use higher rates to control heavy to extremely heavy insect populations.</p> <p>Higher volumes of finished spray may improve insect control under high temperatures when foliage is dense and/or when insect pressure is high.</p> <p>Do not exceed maximum allowable rate.</p> <p>For all listed vegetables types <u>except</u> Root and Tuber Vegetables: Do not apply more than 25.8 fl oz of product or 0.3 pound of active ingredient per acre per season.</p> <p>For Root and Tuber Vegetables: Do not apply more than 21.5 fl oz of product or 0.25 pounds of active ingredient per season.</p>		

Insects Controlled	
<ul style="list-style-type: none"> Aphid spp.¹ Alfalfa Looper Armyworms Cabbage webworm Celery leaf Tier Chinchbug Colorado Potato Beetle Corn Borers Corn Earworm Corn Rootworm (adult) Corn Silkfly Crickets Cucumber beetles Cutworms Diamondback moth Flea beetle Garden Webworm Grasshoppers Ground beetles Hornworms Imported cabbage worm Japanese Beetle (adults) Leafhoppers 	<ul style="list-style-type: none"> Leafminers (adults) Loopers Lygus bugs Meadow Spittlebug Onion Maggot Adults Pepper Maggot (adults) Pepper Weevil Plant bug spp. Potato Leafhopper Saltmarsh Caterpillar Sap Beetle (adults) Southern Corn Rootworm (adult) Stink Bugs Thrips² Tobacco Budworm Tomato Fruitworm Tomato Psyllid Tomato Pinworm Vegetable Weevil Wireworm (adults) Whitefly² Whitefringed Beetle (adult)
<p>¹ Aphid control may be variable depending on species present and host-plant relationships</p> <p>² Aids in control</p>	

Vegetables	Days to Wait Before Harvest	Precautions and Restrictions
Brassica Vegetables Including: Broccoli; Chinese Broccoli (gai lan, white flowering broccoli); Brussels Sprouts; Broccoli Raab (rapini); Cauliflower; Cavalo broccolo; Cabbage; Chinese Cabbage (bok choy); Chinese cabbage (napa); Chinese Mustard Cabbage (gai choy); Collards; Kale; Kohlrabi; Mizuna; Mustard greens; Mustard Spinach; Rape Greens; Turnip Greens	1	Do not make applications less than 7 days apart.
Bulb Vegetables (Allium spp.) Including: Garlic; Great-Headed (elephant); Green Eschalots; Japanese Bunching Onions; Leeks; Onion, Dry Bulb and Green; Onion, Welch; Shallots, Dry Bulb and Green; Spring Onion or Scallions	7	Do not make applications less than 7 days apart. Do not graze livestock in treated areas or cut treated crops for feed.
Fruiting vegetables (except cucurbits) Including: Eggplant; groundcherry (Physalls spp.); Okra; pepino (Melon pear); pepper (includes bell pepper, chili pepper, cooking pepper, pimento, sweet pepper); tomatillo; tomato	1	Do not make applications less than 7 days apart.
Leafy Vegetables (except Brassica) Including: Amaranth (leafy amaranth, Chinese spinach, tampala); Arugula (Roquette); Cardoon; Celery; Chinese; Celtuce; Chervil; Chrysanthemum, edible leaf and garland; Cilantro; (not for use on cilantro grown for seed or coriander); Corn salad; Cress, garden; Cress, upland (yellow rocket, winter cress); Dandelion; Dock (sorrel); Endive (escarole); Fennel, Florence (finocchio); Purslane, winter; Radicchio (red chicory); Rhubarb; Spinach (including New Zealand and vine, Malabar spinach, Indian spinach); Swiss chard	1	Do not make applications less than 7 days apart.
Root and Tuber Vegetables (except Sugar Beet) Including: Arracacha; Arrowroot; Artichoke (Chinese and Jerusalem); Garden Beet; Edible Burdock; Edible Canna; Carrot; Cassava (Bitter and Sweet); Celeriac (Celery Root); Chayote (Root); Turnip-Rooted Chervil; Chicory; Chufa; Dasheen (Taro); Ginger; Ginseng; Horseradish; Leren; Turnip-Rooted Parsley; Parsnip; Potato, Oriental Radish (Daikon); Radish; Rutabaga; Salsify (Oyster Plant); Black Salsify; Spanish Salsify; Skirret; Sweet Potato; Tanier (Cocoyam); Turmeric; Turnip; Yam Bean; and Yam (True)	1	Do not make applications less than 4 days apart. Leaves of Root and Tuber Vegetables cannot be used for food or feed.
Sweet Corn	3	Do not apply within 3 days of harvest. Apply at a minimum of 3 to 5 day intervals or as needed for control.

Precautions and Restrictions:

WARNING: Do not use water based sprays of Zeta-Cyp 1.5 EW OTC Insecticide in conduits, motor housings, junction boxes, switch boxes, or other electrical equipment because of possible shock hazard.

Do not make applications during rain.

Do not water the treated area to the point of run-off.

Do not apply this product in a way that will contact any person or pet, either directly or through drift.

Keep people and pets off treated areas until dry.

Do not treat pets with this product.

Protect outdoor ornamental pools containing fish from spray mist.

For best results, thoroughly wash out sprayer and screen with water and detergent before using Zeta-Cyp 1.5 EW OTC insecticide.

Do not use this product with oil.

All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:

1. Treatment to soil or vegetation around structures;
2. Applications to lawns, turf, and other vegetation;
3. Applications to building foundations up to a maximum height of 3 feet;

Other than applications to building foundations, all outdoor applications to impervious surfaces such as sidewalks, driveways, patios, porches and structural surfaces (such as windows, doors, and eaves) are limited to spot and crack-and-crevice applications only.

Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.

878

Telephone Icon:

For questions or comments call toll-free 1-800-884-3179

Computer Icon:

GordonsUSA.com

[LIMITED WARRANTY AND DISCLAIMER

IMPORTANT: Read this LIMITED WARRANTY AND DISCLAIMER before buying or using this product. By opening and using this product, buyer and all users agree to accept the terms of this LIMITED WARRANTY AND DISCLAIMER in their entirety and without exception. If the terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full.

It is impossible to eliminate all risks inherently associated with use of this product. Damage to the treated article, ineffectiveness, or other unintended consequences can result from use of the product under abnormal conditions such as weather, presence of other materials, or the manner or use of application, etc. Such factors and conditions are beyond the control of the manufacturer, and **BY PURCHASING AND USING THIS PRODUCT THE BUYER AND ALL USERS OF THIS PRODUCT AGREE TO ACCEPT ALL SUCH RISKS.** Buyer and all users further agree to assume all risks of loss or damage from the use of the product in any manner that is not explicitly set forth in or that is inconsistent with label instructions, warnings and cautions.

The manufacturer warrants only that this product conforms to the chemical description given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use, subject to the inherent risks described below. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED.**

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF BUYER AND ALL USERS OF THIS PRODUCT, AND THE EXCLUSIVE LIABILITY OF THE MANUFACTURER, FOR ANY AND ALL LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OF HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OR THE REPAYMENT OF THE PURCHASE PRICE FOR THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THE PRODUCT. The Manufacturer must be promptly notified in writing of any claims, whether based in contract, tort, negligence, strict liability, or otherwise, to be eligible to receive either remedy stated above.

The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere. No employee or agent of the manufacturer or seller is authorized to vary or exceed the terms of this LIMITED WARRANTY AND DISCLAIMER in any manner.]